



## VACCINES REQUIRED FOR CHILD CARE/PRESCHOOL ATTENDANCE July 1, 2012 – June 30, 2013

VACCINE	By 3 Months (on or before last day of mo 2)	By 5 Months (on or before last day of mo 4)	By 7 Months (on or before last day of mo 6)	By 16 Months (on or before last day of mo 15)	By 19 Months (on or before last day of mo 18)
<b>Hepatitis B</b>	<b>1 dose</b> May get dose 1 at birth.	<b>2 doses</b> May get dose 2 as early as 1 month.			<b>3 doses</b>
<b>Diphtheria, Tetanus, Pertussis (DTaP/DT)</b>	<b>1 dose</b>	<b>2 doses</b>	<b>3 doses</b> May get as early as 6 months.		<b>4 doses</b> May get 4 <sup>th</sup> dose as early as 12 months as long as 6 months separate Dose 3 and Dose 4.
<b>Haemophilus influenzae type B (Hib)</b>	<b>1 dose</b>	<b>2 doses</b>	<b>3 doses</b>	<b>4 doses</b>	
<b>Polio (IPV or OPV)</b>	<b>1 dose</b>	<b>2 doses</b> May get as early as 4 months.			<b>3 doses</b>
<b>Pneumococcal Conjugate (PCV7 or PCV13)</b>	<b>1 dose</b>	<b>2 doses</b>	<b>3 doses</b>	<b>4 doses*</b>	
<b>Measles, Mumps, Rubella</b>	<b>Not given before 12 months of age.</b>			<b>1 dose</b>	
<b>Varicella</b>	<b>Not given before 12 months of age.</b>			<b>1 dose</b> OR Healthcare provider verifies disease.	

\*Some children may get 5 total doses. A single supplemental dose of PCV13 is recommended, but not required, for all children aged 14–59 months who got 4 doses of PCV7.

- Refer to *Vaccines Required for School Attendance, Grades K-12* chart for kindergarten entry requirements.
- School-aged children (K-12) in before and after-school programs must meet the immunization requirements for their grade in school.
- Find information on other recommended vaccines not required for child care/preschool attendance: <http://www.immunize.org/cdc/schedules/>

## Minimum Age & Interval for Valid Vaccine Doses

Vaccine	Dose #	Minimum Age	Minimum Interval Between Doses	Notes
Hepatitis B (HepB)	Dose 1	Birth	4 weeks between Dose 1 & 2	<ul style="list-style-type: none"> <li>The final dose in the series should be given at age <math>\geq 24</math> weeks.</li> </ul>
	Dose 2	4 weeks	8 weeks between Dose 2 & 3	
	Dose 3	24 weeks	16 weeks between Dose 1 & 3	
Diphtheria, Tetanus, and Pertussis (DTaP/DT)	Dose 1	6 weeks	4 weeks between Dose 1 & 2	<ul style="list-style-type: none"> <li>Typical vaccine schedule: 2, 4, 6, and 15-18 months of age.</li> <li>Recommended: 6 months between Dose 3 and Dose 4, but <math>\geq 4</math> months is acceptable.</li> </ul>
	Dose 2	10 weeks	4 weeks between Dose 2 & 3	
	Dose 3	14 weeks	6 months between Dose 3 & 4	
	Dose 4	12 months	6 months between Dose 4 & 5	
	Dose 5	4 years	–	
<i>Haemophilus influenzae</i> type B (Hib)	Dose 1	6 weeks	4 weeks between Dose 1 & 2	<ul style="list-style-type: none"> <li>Three doses total for catch-up if 2 doses given before 12 months and Dose 3 given <math>\geq 12</math> months of age. Two doses total for catch-up if Dose 1 given before 12 months and Dose 2 given <math>\geq 12</math> months of age.</li> <li>One dose required if the dose given <math>\geq 15</math> months.</li> <li>If all 3 doses of PedvaxHIB given, only need 3 doses total.</li> </ul>
	Dose 2	10 weeks	4 weeks between Dose 2 & 3	
	Dose 3	14 weeks	8 weeks between Dose 3 & 4	
	Dose 4	12 months	–	
Pneumococcal Conjugate (PCV7 or PCV13)	Dose 1	6 weeks	4 weeks between Dose 1 & 2	<ul style="list-style-type: none"> <li>Three doses total for catch-up if only 2 doses given before 12 months and Dose 3 given <math>\geq 12</math> months of age. Two doses required if both received between 12-24 months at least 8 weeks apart.</li> <li>One dose required if the dose given <math>\geq 24</math> months of age.</li> <li>A single supplemental dose of PCV13 recommended for all children 14–59 months of age who got 4 doses of PCV7.</li> </ul>
	Dose 2	10 weeks	4 weeks between Dose 2 & 3	
	Dose 3	14 weeks	8 weeks between Dose 3 & 4	
	Dose 4	12 months	–	
Polio (IPV or OPV)	Dose 1	6 weeks	4 weeks between Dose 1 & 2	<ul style="list-style-type: none"> <li>Three doses acceptable if child got Dose 3 <math>\geq 4^{\text{th}}</math> birthday.</li> </ul>
	Dose 2	10 weeks	4 weeks between Dose 2 & 3	
	Dose 3	14 weeks	6 months between Dose 3 & 4	
	Dose 4	4 years	–	
Measles, Mumps, and Rubella (MMR or MMRV)	Dose 1	12 months	4 weeks between Dose 1 & 2	<ul style="list-style-type: none"> <li>MMRV (MMR + varicella) may be used in place of separate MMR and varicella vaccines.</li> <li>4-day grace <b>DOES</b> apply between doses of the same live vaccine such as MMR/MMR or MMRV/MMRV. The 4 day grace period <b>DOES NOT</b> apply between Dose 1 and Dose 2 of different live vaccines, such as between MMR and Varicella or between MMR and live flu vaccine.</li> </ul>
	Dose 2	13 months	–	
Varicella (chickenpox) (VAR)	Dose 1	12 months	3 months between Dose 1 & 2	<ul style="list-style-type: none"> <li>Recommended: 3 months between varicella doses, but <math>\geq 28</math> days acceptable.</li> <li>Must get the same day as MMR <b>OR</b> <math>\geq 28</math> days apart (4-day grace <b>DOES NOT</b> apply).</li> </ul>
	Dose 2	15 months	–	

If you have a disability and need this document in another format, please call 1-800-525-0127 (TDD/TTY 1-800-833-6388).